



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

Rec'd PCT/PTO 07 APR 2005
PCT/EP 03/11626

REC'D 24 DEC 2003

WIPO

PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

02023452.2

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk



Anmeldung Nr:
Application no.: 02023452.2
Demande no:

Anmeldetag:
Date of filing: 21.10.02
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

AXXAM S.r.l.
Via Massena, 12/7 int. 1
20145 Milano
ITALIE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Photoprotein with improved bioluminescence

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

C12Q/

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR IE IT LI LU MC NL PT SE SK TR

PHOTOPROTEIN WITH IMPROVED BIOLUMINESCENCE

The present invention provides a chimeric photoprotein with improved luminescent activity, the use thereof as calcium indicator in reporter gene systems and in cell-based assays for the detection and measurement of intracellular calcium.

5 BACKGROUND OF THE INVENTION

Bioluminescence is the capacity of living organisms to emit visible light through a variety of chemiluminescent reaction systems. Bioluminescence reactions require three major components: a luciferin, a luciferase and molecular oxygen. However other components may also be required in some
10 reactions, including cations (Ca^{++} and Mg^{++}) and cofactors (ATP, NAD(P)H). Luciferases are enzymes that catalyse the oxidation of a substrate, luciferin, and produce an unstable intermediate. Light is emitted when the unstable intermediate decays to its ground state, generating oxyluciferin. There are many different unrelated types of luciferin, although many species from at
15 least seven phyla use the same luciferin, known as coelenterazine, which contains a ring formed by three amino acids (2 tyrosines, and a phenylalanine). In some animals (e.g. jellyfish) the luciferin/luciferase system can be extracted in the form of a stable "photoprotein" which emits light upon calcium binding. Photoproteins differ from luciferases in that they are
20 stabilized oxygenated intermediate complexes of luciferase and luciferin. Photoproteins are present in many marine coelenterates and allow these organisms to emit light for a variety of purposes including breeding, feeding, and defense (1). While bacteria emit light continuously, in many other organisms luminescence occurs as flashes, typically of 0.1-1 sec. duration.
25 This requires a rapid switch on/off of the enzymatic reaction and the presence of reagents appropriately sequestered and ready to quick mobilization. In

coelenterates, flashing is caused by calcium entry. The calcium binding sites of photoproteins are homologous to calmodulin. In the presence of calcium, photoproteins emit visible light through an intramolecular reaction. There are many luminescent organisms, but only seven photoproteins, namely

5 Thalassicolin (2,3), Aequorin (4-6), Mitrocromin (syn. Halistaurin) (7,8), Clytin (syn. with Phialidin) (8,9) Obelin (2,6,10,11), Mnemiopsin (12,13), and Berovin (12,13), have been isolated so far. All these proteins are complexes of an apoprotein, an imidazopyrazine chromophore (coelenterazine), and oxygen. Their structures are highly conserved, especially in the region containing the

10 three calcium binding sites (EF-hand structures). These EF-hand structures are characteristic of the calcium-binding protein family. The photoprotein emits light upon reaction with calcium which tightly binds to the EF-hand pocket. The reaction is a single turnover event and results in the release of CO₂ and emission of light in the blue region ($\lambda_{\text{max}}=470$ nm). The term "photoprotein"

15 identifies the luciferin-bound polypeptide, which is capable of luminescence, while "apophotoprotein" is used to indicate the protein without luciferin.

The most studied photoproteins are Aequorin, isolated from *Aequorea victoria* (14) and Obelin, isolated from *Obelia longissima* (15). Upon binding Ca⁺⁺ Aequorin undergoes a conformational change converting itself into an

20 oxygenase (luciferase), which then catalyzes the oxidation of coelenterazine by the bound molecular oxygen. The blue fluorescent protein is made up of coelenteramide, which is an oxidation product of coelenterazine, not covalently bound to apophotoprotein. The photoprotein may be regenerated from the apophotoprotein by incubation with coelenterazine, molecular

25 oxygen, EDTA and 2-mercaptoethanol or dithiothreitol. Since coelenterazine is the common luminescent substrate used by the photoproteins Aequorin, Mitrocromin, Clytin and Obelin, the light-emitting reaction is likely the same in these four photoproteins (16). The recent acquisition of the primary

structure and of the crystallographic data of Aequorin and Obelin gave rise to additional information on their function. Native Obelin from the hydroid *Obelia longissima* is a single-chain protein of 195 amino acid residues (aa) with an approximate molecular mass of 20 kDa that contains the noncovalently bound chromophoric group coelenterazine. The analysis of the primary structures of Clytin shows that it contains 189 aa and belongs to the family of photoproteins. The hydrozoan Ca^{++} -binding photoprotein, however, differ from other Ca^{++} -binding proteins such as calmodulin and troponin C by a relatively high content of cysteine, histidine, tryptophan, proline and tyrosine residues.

Photoproteins are widely used in reporter gene technology to monitor the cellular events associated with signal transduction and gene expression.

The study of cellular events and their regulation requires sensitive, non invasive analytic methods. Photoproteins and in general bioluminescence are excellent reporter systems as they have virtually no background in contrast to fluorescence systems.

Photoproteins have been expressed in mammalian cells to monitor calcium changes in response to different stimuli. Intracellular calcium concentration can be measured by adding coelenterazine cofactor to mammalian cells expressing the photoprotein and detecting photon emission, which is indicative of intracellular calcium concentration.

DESCRIPTION OF THE INVENTION

It has now been found that by chimerization of Obelin protein (Apobelin) through replacement of a region thereof comprised between the first two calcium binding sites, with a corresponding region of a photoprotein selected from Clytin, Aequorin, Thalassicolin, Mitocromin, Mnemiopsoin and Berovin, a novel photoprotein with improved bioluminescence is obtained.

As used herein, Obelin may refer to any of the photoproteins isolated

from different species of *Obelia*, including *Obelia longissima* and *Obelia geniculata* (17). Reference Obelin nucleotide and amino acid sequences are listed in SEQ ID N. 1 and 2, respectively. Reference amino acid and nucleotide sequences of Clytin, Mitocromin and Aequorin are deposited at
5 GenBank accession numbers Q08121, P39047, AAA27720 and, respectively, L13247, L31623, L29571, whereas sequences for Thalassicolin, Mnemiopsis and Berovin are described in references 2, 3, 12 and 13.

A "corresponding region or fragment", as used herein, means any amino acid sequence, within the selected photoprotein, matching Obelin sequence in
10 respective sequence alignments, said region or fragment preferably spanning residues 42-122, more preferably residues 50-95, as referred to Obelin sequence.

According to a preferred embodiment of the invention, the chimeric protein is obtained by replacing residues 50 to 94 of Obelin amino acid
15 sequence with a fragment of Clytin sequence extending from residue 53 to 97. The photoprotein thus obtained, which has been named "Photin", has the amino acid sequence of SEQ ID N. 3.

The chimeric proteins of the invention can be further modified by deletion, addition or substitution of one or more amino acid residues, provided
20 that the activity profile of the photoprotein, in terms of light-emission and calcium-responsiveness, is maintained or increased. Particularly preferred are the substitutions at positions 55, 66, 67, 73, 74, 75, 78, 83, 84, 87, 89 and 94, as referred to obelin sequence.

As shown by *in vitro* studies, Photin produces an intense
25 bioluminescence in response to calcium stimulation, which is generally higher than that observed with natural photoproteins.

For the preparation of the chimeric photoproteins, a recombinant DNA construct bearing portions of the coding sequences of Obelin and of a selected

photoprotein other than Obelin is prepared using conventional genetic engineering, the resulting chimeric product is inserted in a vector, expressed in a suitable host and then isolated and purified. For example, the cDNAs coding for Obelin and for a different photoprotein can be amplified by PCR or constructed *in vitro* with synthetic oligonucleotides and the products can be recombined making use of suitable restriction sites, naturally occurring or artificially introduced into the oligonucleotides used for amplification or for *in vitro* construction. The expression vector can contain, in addition to the recombinant construct, a promoter, a ribosome binding site, an initiation codon, a stop codon or a consensus site for transcription enhancers. The vector can also comprise a selection marker for isolating the host cells containing the DNA construct. Suitable vectors are for example yeast or bacteria plasmids, bacteriophages, viruses, retroviruses or DNA. The vectors carrying the recombinant construct can be introduced in the host by means of conventional techniques. Host cells can be prokaryotic or eukaryotic, such as bacteria, yeasts or mammal cells. The preferred hosts for photoprotein expression/production are mammalian cells, including cells of epithelial or lymphoblast origin such as Hek-293, CHO-K1, HepG2 and HL-60. These cells may express the apophotoprotein in the cytoplasm (18), in the mitochondria, by addition of a mitochondrial targeting sequence to the photoprotein, or in any other cell compartment (19-21). Alternatively, non mammalian cells, such as bacteria or fungi, can be employed. Once the host cell type has been selected, the genetic constructs can be introduced by calcium phosphate precipitation, electroporation or other conventional techniques. After transfection, cells are grown in suitable media and screened for appropriate activities. The photoprotein of the invention may be isolated and purified with conventional procedures, such as extraction, precipitation, chromatography, affinity chromatography, electrophoresis.

Point mutations can be introduced into the chimeric product by means of overlap extension PCR.

To produce Photin, the NdeI/MunI fragment of Obelin gene was replaced with a fragment of 135 nucleotides corresponding to Clytin coding
5 sequence ranging from nucleotide 156 to 291.

In a further aspect, the invention is directed to a nucleic acid molecule encoding the chimeric proteins herein disclosed. DNA coding sequences can be modified by virtue of genetic code degeneracy, for example changing the codon usage to improve the expression in heterologous systems.

10 In a preferred embodiment of the invention, the DNA sequence coding for the Photin protein is selected from SEQ ID No. 4 and 5.

To test whether the chimeric product was functional, in vitro transcription and translation experiments were performed in a cell-free wheat germ translation system in the presence of coelenterazine. In these
15 experiments, active photoprotein formation is recorded by testing luminescence from the translation mixture, after addition of calcium ions. To correlate the measured luminescence with the amount of protein produced, a translation reaction was carried out in the presence of ^{35}S -Methionine. The amount of newly synthesized polypeptide was determined by measurement of
20 ^{35}S -Methionine incorporation into the trichloroacetic insoluble fraction. The results of these experiments demonstrate the sensitivity and effectiveness of Photin in generating bioluminescence in response to calcium-ions stimulation.

In a further embodiment of the invention, the chimeric proteins here provided are used as calcium indicators, especially for the measurement of
25 intracellular calcium concentration. In a typical assay, a cofactor such as coelenterazine is added to mammalian cells expressing the photoprotein and light emission is recorded by methods known in the art, for example using a commercially available luminometer.

Cells expressing both the photoprotein and a receptor involved in intracellular calcium mobilization can be used to test candidate molecules for their effects on receptor modulation. The chimeric photoprotein of the invention can be used in a variety of cell-based functional assays that utilize measurement of intracellular calcium to evaluate the activity of proteins, particularly G-protein coupled receptors (GPCRs) and plasma membrane ion channels. Although the large and rapid increase in intracellular calcium concentration following GPCRs and ion channel stimulation can be detected by various reporters such as calcium-sensitive fluorescent dyes, the use of bioluminescent photoproteins (22) is preferred as their background is virtually absent in contrast to fluorescent dyes; moreover, calcium measurement with photoproteins, besides producing rapid signals, generates a high signal-to-noise ratio with a broad range of detection sensitivity (22, 23). Cell-based functional assays typically comprise adding an appropriate agonist to a culture of cells expressing both GPCRs or ion channels and the photoprotein, and then determining any variation of calcium concentration, for example a calcium increase induced by either rapid influx from extracellular sites or release from intracellular stores (18), through measurements of the photoprotein activity.

The use of cells which express both Photin and a receptor involved in the modulation of intracellular calcium concentration provides a valid system for the screening of compounds for their effects on the release of intracellular calcium. High throughput screening assays can also be designed using a photoprotein according to the invention as reporter system. The sensitivity of the system as well as its high signal to noise ratio allow the use of small assay-volumes.

In a further embodiment, a photoprotein according to the invention is conjugated to a molecule of analytical, diagnostic or therapeutic interest and the conjugation product is used in a competitive solid-phase immunoassay to determine the amount of that molecule in biological samples. For example, the

photoprotein may be chemically conjugated to a hormone protein and used in a solid-phase immunoassay with hormone-specific antibodies to determine the salivary levels of the hormone (24). In a yet further embodiment, a fusion product between a photoprotein according to the invention and a different (poly)peptide, such as a hormone, an antigenic peptide, light or heavy chain immunoglobulin, avidin, streptavidin or protein A, is produced with known techniques of genetic engineering, as disclosed in US6087476 (which is hereby incorporated by reference), and used in immunodiagnostic or imaging processes.

The following examples illustrate the invention in greater detail.

10 DESCRIPTION OF THE DRAWINGS

Fig. 1 Calcium-induced bioluminescence of newly synthesized photoproteins. The photoproteins were translated in a wheat germ cell-free system in the presence of coelenterazine. All translation mixture aliquots were incubated for 2 hrs in the dark at 30°C. The reaction was loaded with CaCl_2 50 mM and luminescence recorded for 10 seconds.

Fig. 2 Photoproteins DNA translation in the presence of coelenterazine, with luminescence measured by diluting the reaction mixture. AQ:Aequorin, PH:Photin, OB:Obelin.

Fig. 3 (A) ATP dose dependent light emission upon stimulation of the endogenous ATP receptor expressed by CHO-K1 cell transfected with photin DNA. Cell transfection, harvesting, and Photin expression were performed as described in Example 1. Cells were treated with two different concentration of ATP. (B) The CHO-K1 cells transfected with Obelin DNA were used as positive control. Luminescence is expressed in RLU (relative light units).

Fig. 4 Dose-dependent curve of Photin activity in CHO-K1 cells transfected with both the Endothelin A receptor and Photin. The receptor was activated by injection of endothelin at a concentration of 100 and 500 nM.

Fig. 5 Dose-dependent curve obtained from cells expressing the

photoprotein Photin and the rat vanilloid receptor 1 (VR1). The specific agonist, capsaicin, was used at a concentration of 10 and 50 μM .

EXAMPLES

1. In Vitro Transcription/Translation of Photoprotein DNA

5 Translation of the photoproteins was carried out in the wheat germ cell-free system (TNT kit, Promega), following the general instructions of the supplier. Approximately 2 μg of DNA was used for each *in vitro* transcription/translation reaction mix. The translation volume (50 μl) included 25 μl of the wheat germ extract, 2 μl of the reaction buffer, a mixture of
10 amino acids except for Methionine, RNasin and Coelenterazine (40 μM), T7 Polimerase. The mixture contained also ^{35}S -Methionine (specific activity 1000 Ci/mmol). ^{35}S -Methionine was used to determine the amount of photoprotein synthesized in vitro. To this end, 5 μl from each sample of the translation mixture were TCA precpitated in ice for 30 min, filtered, washed with cold
15 5% TCA, methanol, dried and placed into counter vials.

The amounts of Photin and Obelin produced in the cell-free system are shown in Table I .

Table I

Inclusion of Radioactive Label into Products of Translation of the photoproteins

Added DNA	Obelin (1 μg)	Photin (1 μg)	Aequorin (1 μg)
^{35}S Counting upon 90 min Translation (cpm)	8169	8290	13049

2. Photoprotein assay

5 and 10 μl of the translation mixture were directly mixed with 95-90 μl
20 of PBS solution in a 96 well plate which was mounted into the Luminometer (Berthold). To trigger photoprotein light emission, 50 μl solution (50 mM CaCl_2) were injected into the well and luminescence recorded for 10 seconds.

The results of the in vitro translation of Photin, Obelin and Aequorin DNA are shown in Fig. 1. The luminescence measurements obtained by diluting the photoprotein in vitro translation reactions are shown in Fig. 2. The comparison of luminescence data with TCA counts obtained during translation with different amounts of Photin and Obelin DNA is shown in Table II (the measured luminescence is proportional to the amount of photoprotein newly synthesized).

Table II

Content of Radiolabeled photoprotein and its Luminescence
in the Translation Mixture

Obelin			Photin	
DNA (ng)	TCA Counts (Cpm/ μ l)	Luminescence (RLU/ μ l)	TCA Counts (cpm/ μ l)	Luminescence (RLU/ μ l)
250	2722	358944	2927	990308
125	2396	271952	2094	858046
63	1030	139241	807	339348
31	579	41970	502	88326
16	265	13482	196	30617

3. Examples of cell-based functional assays

3.1 Photin-expressing clones have been obtained by transfection of CHO-K1 cells (Materials and Methods). Two days after transfection the cells were trypsinized and diluted 10 or 100 times. Once the cells were grown up to well isolated colonies, the colonies were transferred to new plates and selected on the basis of their functional response (luminescent signal) to different concentration of ATP, which is known to stimulate the CHO endogenous receptor and to rise the cytoplasmic Ca^{++} concentration. At the end of each experiment, cells were lysed by perfusion of a solution containing TritonX-100 and CaCl_2 . The active photoprotein was reconstituted incubating the cells with 10 μM coelenterazine diluted in PBS containing 2 mM calcium, in the dark at 37°C in a 5% CO_2 atmosphere for 3 hrs. For light emission measurement, cells were lysed in the presence of calcium and the emitted luminescence recorded. The number of photons emitted during the first 10

seconds was integrated by the luminometer and visualized on the screen. Cells transfected with an empty plasmid or untransfected (data not shown) did not increase photon-emission. To detect changes in calcium concentrations, 10, 50 and 100 μM ATP were injected and the kinetics of calcium response determined. The curve obtained is shown in Fig. 3 (A). A cell line expressing the photoprotein Obelin was used as positive control Fig. 3 (B).

3.2 A CHO cell line expressing endothelin A receptor and Photin was established. Upon stimulation with an agonist, this receptor induces an increase in intracellular calcium concentration which is measured by Photin luminescence. Cells were cultivated as a monolayer in DMEM/F12 medium containing 10% Fetal Bovine Serum (FBS) in a 96-well plate. On the day of the experiment, the culture medium was removed and cells were incubated with 10 μM Coelenterazine in PBS for at least 3 h. Endothelin peptide was diluted in PBS at a concentration of 100 and 500 nM. Approximately 50 μl of endothelin were injected into each well and the response measured. The emitted light was immediately recorded over a period of 30 seconds. The dose-dependent response to endothelin is reported in Figure 4.

3.3 CHO cell lines expressing both Apophotin and the rat VR1 capsaicin receptor - a calcium-permeable ion channel - were grown to 80-95% confluence in tissue culture flasks and harvested by trypsinization. Cells were dispensed at 10,000 cells/well in a 96-well white plates in growth medium and incubated overnight at 37°C in a humidified incubator at 5% CO_2 . For luminescence experiments, the cells were loaded with coelenterazine 10 μM for 3 h at 37°C, 5% CO_2 . Calcium response was stimulated by addition of 10 and 50 μM capsaicin to each well. The kinetics of flash luminescence was followed using Labsystem Luminoskan Ascent, which injects reagents and records light emission from each single well. Addition of capsaicin caused a rapid and transient luminescence signal within 30 seconds, with a peak level

occurring in about 10 seconds (Figure 5).

MATERIALS AND METHODS

Reagents

Restriction enzymes were purchased from New England Biolabs and
5 used according to supplier's instructions. Rnasin and the TNT kit for in vitro
Transcription and Translation were from Promega (Madison, WI). Pfu Turbo
polymerase, reagents for PCR, and competent cells of *E. coli* strains XL-1Blue
and BL21, were from Stratagene (La Jolla, CA). Oligonucleotides were
purchased from Primm (Milan). Coelenterazine was from Prohume Ltd.
10 (Pittsburg, PA). All other chemicals were from standard sources and were of
reagent grade or better.

Assembly PCR for single step synthesis of Obelin DNA sequences

Four steps are required for PCR assembly: oligo synthesis, gene
assembly, gene amplification and cloning. We synthesized 30 oligos, 40 nt in
15 length, which collectively encode both strands of the Obelin gene sequence.
The overlap of complementary oligos is 20 nt. Equal volumes drawn from
each of the 30 oligo solutions were combined to a final concentration of
approximately 250 μ M of mixed oligos, prior to a 250-fold dilution in 20 μ l of
Geneamp XL PCR mix (Perkin Elmer). The amplification process has been
20 carried out in three stages. The first step was carried out with pooled oligos as
follows: 40°C for 2 min, then addition of polymerase, 72°C for 10 sec., then
40 cycles (94°C for 15 s, 40°C for 30 sec. and 72°C for 10 sec.). The reaction
mixture was diluted three-fold with fresh PCR and polymerase mix. The
second stage was: 25 cycles (94°C for 15 s, 40°C for 30 s and 72°C for 45 s).
25 The reaction was again diluted threefold in complete PCR mix. The conditions
for the third stage were: 20 cycles (94°C for 15 s, 40°C for 30 s and 72 °C for
70 s). The reaction products were analyzed by 1% agarose-gel electrophoresis,
and the specific fragment cloned into the pcrBlunt vector for further cloning

steps. The plasmid has been called pcr-OB. The fidelity of the PCR reactions was confirmed by dideoxy sequencing.

Construction of the chimeric protein

Four couples of oligos have been designed from the Clytin gene sequence. The region between EF-hand I and EF-hand II have been chosen for gene chimerization. The oligonucleotide primers were the following:

Clyt-4 5'

AATTCTTTCCATCCATCAACAAAAGCTGGGAATTCGACTTCTTTACC
ATAATCCATACCAATC-3'

10 Clyt-3 5'-

AAAGATTGGTATGGATTATGGTAAAGAAGTCGAATTCCCAGCTTTTG
TTGATGGATGGAAAG-3'

Clyt-2 5'-

TTTTTGAAGAAAGCTTCGACAGCATCCTGGTGACGTTTGGTCTGTTCT
15 GGTGTTGCTCCAAGTTTGGCGCA-3'

Clyt-1 5'-

TATGCGCCAAACTTGGAGCAACACCAGAACAGACCAAACGTCACCA
GGATGCTGTCGAAGCTTTCTTCAA-3'

The annealed oligos have been cloned into the *NdeI/MunI* unique sites in the pcr-OB vector. The plasmid containing the chimeric gene product is called pcr-Photin. Photin DNA has been further subcloned into the poDNA3 vector which contains the T7 promoter.

PCR-Based Codon Usage Change

The technique of overlap extension PCR was employed to produce the Obelin mutant. Six couples of primers were designed with ten different point mutations which produce ten different codon usage changes. To amplify DNA fragments used to overlap extension, PCR reactions were carried out with 2.5 units of Pfu polymerase, 50 ng of DNA template, 250 μ M each dNTP, and 50

pmol of each primer in a total volume of 100 μ l. The cycling parameters employed were 94°C for 1 min, 45° for 1 min, and 72°C for 1 min 30 s for the first 10 cycles, followed by 20 cycles with an annealing temperature of 50°C. The site-specific mutations were confirmed through DNA sequencing
5 performed at Primm (Milan). All molecular procedures were conducted using standard protocols.

CHO Cell Culture

All cells were cultured under standard humidified conditions at 37°C and 5% CO₂. CHO cells were maintained in DMEM/F12 + FBS 10% +
10 Pen/Strep + G418 0.5 mg/ml + Pyruvate 1.6 mM + NaHCO₃ 0.2%, all reagents were from Life Technologies. DNA transfection was performed when these cells were grown to a 70% to 80% confluence on the plates.

REFERENCES

1. Kendall, J. M., and Badminton, M.N. (1998). *Aequorea victoria*. bioluminescence moves into an exciting new era. Trends Biotechnology. 16(5):216-24.
- 5 2. Campbell, A. K., Hallet, R.A., Daw, M.E., Ryall, R.C., Hart and Herring P.J. (1981). Application of the photoprotein obelin to the measurement of free Ca^{++} in cells. In Bioluminescence and Chemiluminescence, basic Chemistry and Analytical applications (Edited by M.A. deLuca and W. D. McElroy), pp. 601-607. Academy Press, New York.
- 10 3. Herring, P.J. (1979) Some features of the bioluminescence of the radiolarian *Thalassicola sp.* Mar. Biol. 53, 213-216.
4. Shimomura, O., Johnson F.H., and Saiga, Y (1962) Extraction, purification and properties of aequorin, a bioluminescent protein from the luminous hydromedusan, *Aequorea*. J.Cell.Comp.Physiol. 59,223-239.
- 15 5. Shimomura, O., Johnson F.H., and Saiga, Y (1963) Further data on the bioluminescent protein, aequorin. J.Cell.Comp.Physiol. 62, 1-8.
6. Morin, J.G. and Hastings J.W. (1971) Biochemistry of the bioluminescence of colonial hydroids and other coelenterates. J.Cell. Physiol. 77, 305-311.
- 20 7. Shimomura, O., Johnson, F.H. and Saiga, Y. (1963) Extraction and properties of halistaurin, a bioluminescent protein from the hydromedusan *Halistaura*. J.Cell. Physiol. 62, 9-15.
8. Shimomura, O., and Shimomura, A. (1985) Halistaurin, phialidin and modified forms of aequorin as Ca^{++} indicator in biological systems. Biochem. J. 228,745-749.
- 25 9. Levine, L.D., and Ward, W.W. (1982) Isolation and characterization of a photoprotein "phialidin" and a spectrally unique green-fluorescent protein

from the bioluminescent jellyfish *Phialidium gregarium*. Comp. Biochem. Physiol. 72B, 77-85.

10. Morin, J.G. and Hastings (1971) Energy transfer in a bioluminescent system. J. Cell. Physiol. 77, 313-318.

5 11. Campbell, A.K. (1974) Extraction, partial purification and properties of obelin the calcium-activated protein from the hydroid *Obelia geniculata*. Biochem. J. 143, 411-418.

12. Ward, W.W. and Selinger (1974) Extraction and purification of calcium-activated photoprotein from the ctenophores *Mnemiopsis sp.* and *Beroë ovata*. Biochemistry 13, 1491-1499.

13. Ward, W.W. and Selinger H.H. (1974) Properties of mnemiopsin, and berovin, calcium-activated photoproteins from the ctenophores *Mnemiopsis sp.* and *Beroë ovata*. Biochemistry 13, 1500-1510.

14. Johnson, F.H. and Shimomura, O. (1978) Introduction to the bioluminescence of medusae, with special reference to the photoprotein aequorin. Methods Enzymol. 57, 271-291.

15. Illarionov BA, Bondar VS, Illarionova VA, Vysotski ES. Sequence of the cDNA encoding the Ca^{++} -activated photoprotein obelin from the hydroid polyp *Obelia longissima*. Gene. 1995 14;153(2):273-4.

20 16. Blinks, J.R., Weir, W.G., Hess, P. and Prendergast, F.G. (1982). Measurement of Ca^{++} concentrations in living cells. Prog. Biophys. Mol. Biol. 40, 1-114.

17. Markova SV, Vysotski ES, Blinks JR, Burakova LP, Wang BC, Lee J., (2002) Obelin from the bioluminescent marine hydroid *Obelia geniculata*: cloning, expression, and comparison of some properties with those of other Ca^{2+} -regulated photoproteins. Biochemistry. 2002 Feb 19; 41(7):2227-36.

25 18. Button D, Brownstein M. (1993) Aequorin-expressing mammalian cell lines used to report Ca^{++} mobilization. Cell Calcium. Oct; 14(9):663-71.

19. Mattheakis, L., and Ohler, L.D. (2002) Seeing the light: Calcium imaging in cells for drug discovery. *Drug Discovery Today S*, 15-19.
20. Stables J, Green A, Marshall F, Fraser N, Knight E, Sautel M, Milligan G, Lee M, Rees S. (1997). A bioluminescent assay for agonist activity at potentially any G-protein-coupled receptor. (1997) *Anal Biochem* 1;252(1):115-26.
21. Brini M, Pinton P, Pozzan T, Rizzuto R. (1999). Targeted recombinant aequorins: tools for monitoring Ca^{++} in the various compartments of a living cell. *Microsc.Res.Tech.* 46,380-389.
- 10 22. Creton R, Kreiling JA, Jaffe LF. (1999) Calcium imaging with chemiluminescence. *Microsc.Res. Tech.* 46,390-397.
23. Shimomura, O. et al. (1993) Light-emitting properties of recombinant semi-synthetic Aequorins and recombinant fluorescein-conjugated Aequorin for measuring cellular calcium. *Cell Calcium* 14, 373-378.
- 15 24. Mirasoli, M. et al. (2002) Bioluminescence immunoassay for cortisol using recombinant aequorin as a label. *Analytical Biochemistry* 306, 204-211.

SEQUENCE LISTING

<110> Axxam

5 <120> photoprotein with increased bioluminescent activity

<130> 1191EUR

<160> 9

10

<170> PatentIn version 3.1

<210> 1

<211> 662

15

<212> DNA

<213> Obelia longissima

<400> 1

20

acgatcgaac caaacaactc agotcacagc tactgaacaa ctcttgttgt gtacaatcaa 60

aatgtcttca aaatacgcag ttaaactcaa gactgacttt gataatccac gatggatcaa 120

aagacacaag cacatgtttg atttcctcga oatcaatgga aatggaaaaa tcaccctoga 180

25

tgaaattgtg tccaaggcat ctgatgacat atgtgccaaag ctggaagcca caccagaaca 240

aagaaaaacgc catcaagttt gtgttgaagc tttctttaga ggatgtggaa tggaatatgg 300

30

taaagaaatt gccttcccac aattcctcga tggatggaaa caattggcga cttcagaact 360

caagaaatgg gcaagaaacg aacctactct cattcgtgaa tggggagatg ctgtctttga 420

tatttttcgac aaagatggaa gtggtacaat cactttggac gaatggaaag cttatggaaa 480

35

aatctctggg atctctccat cacaagaaga ttgtgaagcg acatttcgac attgcgattt 540

ggacaacagt ggtgaccttg atgttgaagc gatgacaaga caacatcttg gattctggta 600

40

cactttggac ccagaagctg atggtctcta tggcaacgga gttccctaag othtttttcg 660

aa 662

<210> 2

45

<211> 195

<212> PRT

<213> Obelia longissima

<400> 2

50

Met Ser Ser Lys Tyr Ala Val Lys Leu Lys Thr Asp Phe Asp Asn Pro
1 5 10 15

55

Arg Trp Ile Lys Arg His Lys His Met Phe Asp Phe Leu Asp Ile Asn
20 25 30

60

Gly Asn Gly Lys Ile Thr Leu Asp Glu Ile Val Ser Lys Ala Ser Asp
35 40 45

5	Asp	Ile	Cys	Ala	Lys	Leu	Glu	Ala	Thr	Pro	Glu	Gln	Thr	Lys	Arg	His
	50						55					60				
10	Gln	Val	Cys	Val	Glu	Ala	Phe	Phe	Arg	Gly	Cys	Gly	Met	Glu	Tyr	Gly
	65					70					75					80
15	Lys	Glu	Ile	Ala	Phe	Pro	Gln	Phe	Leu	Asp	Gly	Trp	Lys	Gln	Leu	Ala
					85					90					95	
20	Thr	Ser	Glu	Leu	Lys	Lys	Trp	Ala	Arg	Asn	Glu	Pro	Thr	Leu	Ile	Arg
				100					105					110		
25	Glu	Trp	Gly	Asp	Ala	Val	Phe	Asp	Ile	Phe	Asp	Lys	Asp	Gly	Ser	Gly
			115					120					125			
30	Thr	Ile	Thr	Leu	Asp	Glu	Trp	Lys	Ala	Tyr	Gly	Lys	Ile	Ser	Gly	Ile
		130					135					140				
35	Ser	Pro	Ser	Gln	Glu	Asp	Cys	Glu	Ala	Thr	Phe	Arg	His	Cys	Asp	Leu
	145					150					155					160
40	Asp	Asn	Ser	Gly	Asp	Leu	Asp	Val	Asp	Glu	Met	Thr	Arg	Gln	His	Leu
					165					170					175	
45	Gly	Phe	Trp	Tyr	Thr	Leu	Asp	Pro	Glu	Ala	Asp	Gly	Leu	Tyr	Gly	Asn
				180					185					190		
50	Gly	Val	Pro													
		195														
55	<210>	3														
	<211>	195														
	<212>	PRT														
	<213>	chimeric protein														
60	<400>	3														
65	Met	Ser	Ser	Lys	Tyr	Ala	Val	Lys	Leu	Lys	Thr	Asp	Phe	Asp	Asn	Pro
	1				5					10					15	
70	Arg	Trp	Ile	Lys	Arg	His	Lys	His	Met	Phe	Asp	Phe	Leu	Asp	Ile	Asn
				20					25					30		
75	Gly	Asn	Gly	Lys	Ile	Thr	Leu	Asp	Glu	Ile	Val	Ser	Lys	Ala	Ser	Asp
		35						40					45			
80	Asp	Ile	Cys	Ala	Lys	Leu	Gly	Ala	Thr	Pro	Glu	Gln	Thr	Lys	Arg	His

20

50

55

60

5 Gln Asp Ala Val Glu Ala Phe Phe Lys Lys Ile Gly Met Asp Tyr Gly
65 70 75 80

10 Lys Glu Val Glu Phe Pro Ala Phe Val Asp Gly Trp Lys Glu Leu Ala
85 90 95

Thr Ser Glu Leu Lys Lys Trp Ala Arg Asn Glu Pro Thr Leu Ile Arg
100 105 110

15 Glu Trp Gly Asp Ala Val Phe Asp Ile Phe Asp Lys Asp Gly Ser Gly
115 120 125

20 Thr Ile Thr Leu Asp Glu Trp Lys Ala Tyr Gly Lys Ile Ser Gly Ile
130 135 140

25 Ser Pro Ser Gln Glu Asp Cys Glu Ala Thr Phe Arg His Cys Asp Leu
145 150 155 160

30 Asp Asn Ser Gly Asp Leu Asp Val Asp Glu Met Thr Arg Gln His Leu
165 170 175

Gly Phe Trp Tyr Thr Leu Asp Pro Glu Ala Asp Gly Leu Tyr Gly Asn
180 185 190

35 Gly Val Pro
195

40 <210> 4
<211> 600
<212> DNA
<213> chimeric gene

45 <400> 4
ggatccgccg ccatgtcttc aaaatacgca gttaaactca agactgaatt tgataatcca 60
cgatggatca aaagacacaa gcacatgttt gatttctctg acatcaatgg aaatggaaaa 120
50 atcaccctcg atgaaattgt gtccaaggca tctgatgaca tatgcgcgcaa acttggagca 180
acaccagaac agaccaaacg tcaccaggat gctgtcgaag atttcttcaa aaagattggt 240
55 atggattatg gtaaagaagt cgaattccca gcttttgttg atggatggaa agaattggcg 300
acttcagaac tcaagaaatg ggcaagaaac gaacctactc tcattcgtga atggggagat 360
gctgtctttg atattttcga caaagatgga agtggtacaa tcacttttga cgaatggaaa 420
60 gcttatggaa aaatctotgg tatctctcca tcacaagaag attgtgaagc gacatttcga 480
cattgcgatt tggacaacag tgggtgacctt gatgttgacg agatgacaag acaacatctt 540

21

ggattctggt acactttgga cccagaagct gatggtctct atggcaacgg agttccctaa 600

5 <210> 5
 <211> 607
 <212> DNA
 <213> chimeric gene

10 <400> 5
 ggatccgccc coactgtcttc aaaatacgcg gttaaactca agactgactt tgataatcca 60

cgatggatca aaagacacaa gcacatgttt gatttcctcg acatcaatgg aaatggaaaa 120

15 atcaccctcg atgaaattgt gtccaaggca tetgatgaca tctgcgcaca actgggagca 180

acaccagaac agaccaaaacg gcaccaggat gctgtcgaag ctttcttcaa aaagattggt 240

20 atggattatg gtaaagaagt cgaattccca gcttttgttg atggatggaa agaattggcg 300

acttcagaac tcaagaaatg ggcaagaaac gaacctactc tcattogtga atggggagat 360

gctgtctttg atattttcga caaagatgga agtgggtacaa tcactttgga cgaatggaaa 420

25 gcttatggaa aaatctctgg tatctotcca tcacaagaag attgtgaagc gacatttcga 480

cattgcatc tggacaacag tggcgacctg gatgttgacg agatgacaag acaacatctt 540

30 ggattctggt acactttgga cccagaagct gatggcctct atggcaacgg agttccctaa 600

gaattcc 607

35 <210> 6
 <211> 70
 <212> DNA
 <213> oligonucleotide primer

40 <400> 6
 tatgcgccaa acttggagca acaccagaac agaccaaaacg tcaccaggat gctgtcgaag 60
 ctttcttcaa 70

45 <210> 7
 <211> 71
 <212> DNA
 <213> oligonucleotide primer

50 <400> 7
 tttttgaaga aagcttogac agcatcctgg tgacgtttgg tctgttctgg tgttgctcca 60
 agtttggcgc a 71

55 <210> 8
 <211> 62
 <212> DNA
 <213> oligonucleotide primer

60 <400> 8
 aaagattggt atggattatg gtaaagaagt cgaattccca gcttttgttg atggatggaa 60

22

ag 62

5 <210> 9
<211> 63
<212> DNA
<213> oligonucleotide primer

10 <400> 9
aattctttcc atccatcaac aaaagctggg aattogactt cttaccata atccatacca 60
ato 63

CLAIMS

1. A chimeric photoprotein obtained by replacing a region of Obelin protein comprised between the first and the second calcium binding sites with a corresponding region of a photoprotein selected from Clytin, Aequorin,
5 Thalassicolin, Mitocromin, Mnemiopsin and Berovin.
2. A chimeric photoprotein according to claim 1, wherein said region is from residue 42 to 122 of Obelin protein sequence.
3. A chimeric photoprotein according to claim 2, wherein said region extends from residue 50 to 95 of Obelin protein sequence.
- 10 4. A chimeric photoprotein according to claims 1-3, in which residues 50 to 94 of Obelin protein are replaced with a fragment of Clytin sequence extending from residue 53 to 97.
5. A chimeric photoprotein according to claim 4, having the amino acid sequence of SEQ ID N. 3.
- 15 6. A chimeric photoprotein according to claims 4-5, further comprising one or more amino acid substitutions at positions 55, 66, 67, 73, 74, 75, 78, 83, 84, 87, 89 and 94 of Obelin sequence.
7. A fusion product between a photoprotein according to claims 1-6 and a different (poly)peptide.
- 20 8. A conjugation product between a photoprotein according to claims 1-6 and a molecule of analytical, diagnostic or therapeutic interest.
9. An isolated nucleic acid molecule encoding the protein of claim 3, having a sequence selected from SEQ. ID N. 4 and SEQ ID N. 5.
10. The use of a chimeric photoprotein according to claims 1-6, in
25 combination with a luciferin substrate, for the detection of calcium ions.
11. The use according to claim 10, wherein said luciferin substrate is coelenterazine.

12. The use according to claims 10-11, for the quantitative determination of calcium ions.
13. The use according to claims 10-11, for the determination of intracellular calcium concentration.
- 5 14. A host cell bearing a nucleic acid molecule according to claim 9.
15. The cellular host of claim 14, which is selected from bacterial, yeast, fungal, plant, insect and animal cells.
16. A method for producing Photin protein, which comprises growing the recombinant host cell of claim 14 in conditions suitable for photoprotein
10 expression, and recovering the expressed protein.
17. A method for the screening of biologically active molecules, which comprises exposing a cellular host according to claim 14 to a definite amount of said molecules and detecting any variation of intracellular calcium concentration.
- 15 18. A method according to claim 17, wherein the host cell is transfected with a heterologous G-protein coupled receptor or ion channel.
19. The use of a conjugation product between a photoprotein according to claims 1-6 and a molecule of analytical, diagnostic or therapeutic interest in a competitive solid-phase immunoassay for determining the amount of said
20 molecule in biological samples.
20. A Bioluminescence resonance energy transfer (BRET) system, comprising a fluorescent protein and the photoprotein of claim 5.

25

ABSTRACT**PHOTOPROTEIN WITH IMPROVED BIOLUMINESCENCE**

- 5 Disclosed is a photoprotein with improved luminescent activity, the use thereof as calcium indicator and in cell-based assays for the detection and measurement of intracellular calcium.

SHEET 1/5




FIGURE 1

SHEET 2/5





FIGURE 2

SHEET 3/5

(A)



(B)




FIGURE 3

SHEET 4/5




FIGURE 4

SHEET 5/5




FIGURE 5